

Association between an anti-inflammatory low-carbohydrate diet and gingivitis in women with obesity

Khadiga O Ali¹, Mohammad M Hammad¹, Rola A Al-Habashneh¹, Jowan M Al-Nusair², Mustafa M Ababneh³, Mohammad B Al-Zghoul¹

ABSTRACT

Background: The Western diet contributes to the onset and progression of gingivitis. In contrast, an anti-inflammatory low-carbohydrate diet (AILCD) has been proposed as a dietary recommendation to improve gingival health. Although a few published studies have investigated the influence of an anti-inflammatory diet on gingivitis in individuals of normal weight, this study aimed to evaluate the association between an AILCD and gingivitis in women with obesity. **Methods:** Twenty-three women with obesity and gingivitis participated in the 2-month study. Clinical parameters, including plaque index (PI), gingival index (GI), probing pocket depth (PPD), and bleeding on probing (BOP), were recorded at baseline, 1 month, and 2 months. Mouthwash samples were collected for real-time PCR detection of oral microbiota, including *Streptococcus mutans*, *Aggregatibacter actinomycetemcomitans*, *Tannerella forsythia*, *Porphyromonas gingivalis*, and *Fusobacterium nucleatum*. Salivary inflammatory cytokines (IL-6, TNF- α , and IL-10) were measured using ELISA. Anthropometric parameters, including weight, body mass index (BMI), skeletal muscle mass (SMM), and percent body fat (PBF), were assessed using body composition analysis. **Results:** After 2 months, all clinical parameters were significantly reduced. The baseline PI was 1.14 ± 0.48 reduced to 0.54 ± 0.38 , GI was 0.77 ± 0.32 reduced to 0.38 ± 0.17 , PPD was $1.83 \text{ mm} \pm 0.22 \text{ mm}$ reduced to $1.35 \text{ mm} \pm 0.30 \text{ mm}$, and BOP was reduced from 37% to 19%. Baseline weight, BMI, SMM, and PBF were significantly reduced by 3.8 kg, 1.3 kg/m², 0.5 kg, and 1.6%, respectively. However, no significant differences were observed in the periodontal pathogens and salivary inflammatory cytokines except for *Tannerella forsythia*, which showed a significant increase in its counts with a *p* value of 0.01. **Conclusion:** The AILCD followed by participants in this study was associated with significant improvements in clinical and anthropometric parameters. However, *Tannerella forsythia* showed a significant increase in women with obesity.

RÉSUMÉ

Contexte : Le régime alimentaire occidental contribue à l'apparition et à la progression de la gingivite. En revanche, un régime anti-inflammatoire à faible teneur en glucides (AILCD) a été proposé comme recommandation alimentaire pour améliorer la santé gingivale. Bien que quelques études publiées aient étudié l'influence d'un régime anti-inflammatoire sur la gingivite chez les personnes de poids normal, la présente étude visait à évaluer l'association entre un régime AILCD et la gingivite chez les femmes souffrant d'obésité. **Méthodes :** Vingt-trois femmes souffrant d'obésité et de gingivite ont participé à l'étude de 2 mois. Les paramètres cliniques, notamment l'indice de plaque (PI), l'indice gingival (GI), la profondeur de la poche parodontale (PPP) et le saignement au sondage (BOP), ont été enregistrés au départ, à 1 mois et à 2 mois. Des échantillons de rince-bouche ont été prélevés pour analyse PCR en temps réel du microbiote oral, notamment à la recherche de *Streptococcus mutans*, *Aggregatibacter actinomycetemcomitans*, *Tannerella forsythia*, *Porphyromonas gingivalis* et *Fusobacterium nucleatum*. Les cytokines salivaires inflammatoires (IL-6, TNF- α et IL-10) ont été mesurées à l'aide d'essais ELISA. Les paramètres anthropométriques, notamment le poids, l'indice de masse corporelle (IMC), la masse des muscles squelettiques (MMS) et le pourcentage de graisse corporelle (PGC), ont été évalués au moyen d'une analyse de la composition corporelle. **Résultats :** Après 2 mois, tous les paramètres cliniques étaient considérablement réduits. L'index PI au départ (base référence) était de $1,14 \pm 0,48$ ramené à $0,54 \pm 0,38$, l'index GI était de $0,77 \pm 0,32$ ramené à $0,38 \pm 0,17$, la PPP était de $1,83 \text{ mm} \pm 0,22 \text{ mm}$ ramenée à $1,35 \text{ mm} \pm 0,30 \text{ mm}$, et la mesure BOP était réduite de 37 % à 19 %. La pondération, l'IMC, la MMS et le PGC au départ (base référence) ont été considérablement réduits de 3,8 kg, 1,3 kg/m², 0,5 kg et 1,6 %, respectivement. Toutefois, aucune différence significative n'a été observée dans les pathogènes parodontaux et les cytokines salivaires inflammatoires, à l'exception de *Tannerella forsythia*, dont on a constaté une augmentation significative des dénombrements avec une valeur *p* de 0,01. **Conclusion :** Le régime alimentaire AILCD suivi par les participants à l'étude a été associé à des améliorations significatives des paramètres cliniques et anthropométriques. Toutefois, la présence de *Tannerella forsythia* est associée à une augmentation significative du nombre de femmes souffrant d'obésité.

Keywords: anti-inflammatory low-carbohydrate diet; body mass index; gingivitis; obesity; periodontal pathogens
CDHA Research Agenda category: risk assessment and management

¹Department of Preventive Dentistry, Faculty of Dentistry, Jordan University of Science & Technology, Irbid, Jordan

²Department of Internal Medicine, Joan C. Edwards School of Medicine, Marshall University, Huntington, WV, USA

³Department of Basic Veterinary Medical Sciences, Faculty of Veterinary Medicine, Jordan University of Science and Technology, Irbid, Jordan

Correspondence: Prof. Rola Al-Habashneh; email: raalhabashneh@just.edu.jo

Manuscript submitted 27 July 2024; revised 3 September and 24 December 2024, 22 February 2025; accepted 11 April 2025

©2026 Canadian Dental Hygienists Association

PRACTICAL IMPLICATIONS OF THIS RESEARCH

- Evidence of the association between anti-inflammatory low-carbohydrate diets (AILCD) and gingivitis among women with obesity is limited.
- This study showed that an AILCD was associated with improved clinical and anthropometric parameters. However, *Tannerella forsythia* increased significantly in women with obesity during this study duration.
- AILCDs may lead to weight loss and improve gingival health in women with obesity, but further research is indicated.

INTRODUCTION

Biofilm-induced gingivitis (BIG) is a site-specific inflammatory condition caused by dental biofilm deposition, characterized by gingival inflammation without attachment loss or alveolar bone loss.^{1,2} Epidemiological studies indicate that BIG is more common and severe in individuals who neglect oral hygiene.³ The severity of periodontal diseases ranges from reversible gingivitis to more severe, irreversible periodontitis, which is considered a risk factor for long-term tooth loss.⁴ Moreover, periodontitis is associated with more serious systemic conditions such as diabetes⁵ and cardiovascular disease (CVD).⁶

The literature provides substantial evidence that the accumulation of dental biofilm is the primary cause of gingivitis. However, its role in promoting gingival inflammation is mainly observed in Western diet environments.^{1,7} Dental biofilm is composed of a complex community of bacteria that accumulate along the gingival margin and involves the initial colonization by gram-positive bacteria, such as *Streptococcus mutans*,⁸ a member of the “oral streptococci” group. *S. mutans* produces sticky, branched glucose polymers known as glucans, which play a critical role in the formation and adhesion of dental plaque by providing a matrix for bacterial attachment and biofilm development. In addition to glucans, other components such as proteins, lipids, and extracellular polysaccharides also contribute to plaque formation. When dysbiosis exceeds the immune system’s capacity to control it, inflammation extends subgingivally, increasing the proportion of pathogenic gram-negative anaerobic bacteria and driving the progression of periodontitis.^{9,10}

The relationships between different bacterial species in dental plaque are complex. The presence, absence, and relative proportions of these species are influenced by various factors, including the host’s diet.¹¹ Therefore, dietary recommendations can help manage gingival and periodontal inflammation,¹²⁻¹⁴ as well as systemic chronic diseases such as diabetes and CVD.¹³ Omega-3 fatty acids in fatty fish reduce inflammation in both the gingiva and cardiovascular system.¹⁵ Antioxidants, such as vitamins C and E from fruits and vegetables, improve gingival health by neutralizing free radicals.¹⁶ Additionally, limiting processed sugars and refined carbohydrates helps regulate blood glucose and inflammation, benefiting both periodontal and systemic health.¹⁷

The role of a highly refined carbohydrate diet in supragingival plaque accumulation and dental caries is well known.¹⁸ Additionally, it has been reported that a reduction in carbohydrate intake reduces gingivitis despite constant plaque values.^{7,19} In a study lasting 4 weeks, an anti-inflammatory diet (AID) reduced the load of the oral Streptococci group, *Granulicatella adiacens*, *Actinomyces* spp., and *Fusobacterium* spp. in supragingival plaque.²⁰

El Makaky et al.²¹ studied the effects of a 4-week

AID that was low in processed carbohydrates and high in vitamin C, antioxidants, omega-3 fatty acids, plant nitrates, vitamin D, and fibre on clinical and serological inflammatory markers in children with gingivitis. They concluded that an AID can significantly reduce serological markers and gingival inflammation in children. Woelber et al.²² studied the influence of an AID on gingivitis in a randomized controlled trial. They observed that this diet could significantly reduce gingivitis in normal-weight individuals, although serological inflammatory markers and the subgingival microbiota appeared to be unaltered during the 4-week course of the study. Moreover, Rajaram et al.¹² studied the effect of an AID on gingival inflammatory parameters and found that this type of diet significantly reduces gingival inflammation in individuals with chronic gingivitis.

Furthermore, studies have shown a correlation between obesity and the severity of periodontal disease; the majority of the meta-analyses performed revealed that individuals with obesity had significantly greater clinical parameters of gingival inflammation.^{23,24} An association has also been identified between obesity and periodontal disease in women, with the connection becoming more pronounced as obesity levels rise.^{25,26} Interestingly, this pattern is not observed in men with an elevated body mass index (BMI). This difference may be attributed to the fact that, in men, a higher BMI often reflects greater muscle mass or bone density rather than body fat. In addition, the distribution of body fat varies between genders, so a higher BMI in men does not necessarily indicate obesity.²³

No known studies in the literature have investigated the association between anti-inflammatory low-carbohydrate diets (AILCD) and gingivitis in women with obesity. Therefore, this study aimed to evaluate the association between AILCD and clinical parameters, periodontal pathogens, and salivary inflammatory cytokines in women with obesity and gingivitis using real-time polymerase chain reaction (RT-PCR) over a 2-month duration to investigate a dietary intervention as an adjunctive method for the resolution of gingivitis.

Study hypothesis

This study hypothesized that AILCD would reduce gingival inflammation, improve clinical and anthropometric parameters, and lead to positive changes in salivary inflammatory cytokines and periodontal pathogens in women with obesity and gingivitis after 2 months, compared to baseline.

MATERIALS AND METHODS

Study design

The Institutional Review Board (IRB) of Jordan University of Science and Technology (JUST) approved this non-controlled repeated measures clinical trial (Reference

number: 196/2022). Additionally, the study was registered for funding under Grant number 20220426. It was conducted at the Preventive Dentistry Department (Periodontics Clinics) of Jordan University of Science and Technology, Irbid, Jordan, between October 2022 and July 2023. Participants were informed of the study's objectives and methodology, and written informed consent was obtained.

Sample selection

Women selected for the study fulfilled the following criteria:

1. Generalized gingivitis (BOP $\geq 10\%$)²⁷
2. Western diet conditions with a processed carbohydrate intake $>45\%$ ²⁸
3. BMI ≥ 30 kg/m²
4. Age 18 to 45 years

The exclusion criteria for the study were as follows:

1. Smoking
2. Severe or life-threatening illnesses
3. Intake of antibiotics within 3 months before the start of or during the study period
4. Drugs influencing gingival inflammation or bleeding (e.g., anticoagulants, cortisone)
5. Carbohydrate- or insulin-related diseases (e.g., diabetes)
6. Pregnancy or breastfeeding

Recruitment was based on a Google questionnaire, which included specific questions to determine whether potential participants met the inclusion criteria, along with advertisements at Jordan University of Science and Technology, the JUST Dental Center, and a private nutrition centre.

Dietary intervention

All participants who met the inclusion criteria underwent testing for vitamin D and C levels, and then were instructed to follow an AILCD under the supervision of the same dietitian for 2 months, with in-person follow-up visits every 10 days. During these visits, participants engaged in a 1-hour discussion with the dietitian to evaluate their progress, adherence, and any challenges faced during the previous period. Compliance was monitored using written dietary checklists. Frequent follow-up visits were designed to ensure participants' adherence to the diet and provide periodic motivation. Participants were instructed to maintain their routine oral hygiene methods and avoid changes to their physical activity during the observation period.

The dietary recommendations were largely based on those of a study by Woelber et al.,¹⁹ with some adjustments that omega-3 was not taken as a supplement but rather was obtained through the diet. The diet was a 1500-calorie program with 20% carbohydrates (75 g/d), 25% protein (94 g/d), and 55% fat (92 g/d). Participants had 3 main meals per day without snacks. The following components were included in the dietary intervention:

Macronutrients

- Omission of processed carbohydrates such as sugar, honey, white flour, white rice or fruit juices; reduction of the intake of starches, as far as possible, to 75 g/day.²⁸
- No restrictions on fruits and vegetables (polysaccharides).
- Intake of omega-3 fatty acids (salmon, walnuts, and beans), a restriction on the amount of transfatty acids as far as possible (such as fried meals, crisps, doughnuts, and croissants), and a reduction in omega-6 fatty acids as far as possible (such as safflower oil, grape seed oil, sunflower oil, margarine, sesame oil, and corn oil).
- Reduction of processed animal proteins (such as sausages and cold cuts) as far as possible, favouring plant proteins (such as legumes and nuts).

Micronutrients

- Daily intake of a source of vitamin C (such as 2 kiwis, 1 orange, 1 bell pepper).
- Daily intake of a source of vitamin D either by exposing the body to 15 min unprotected in the sun or supplementation with 1,000 international units (25 µg). In the case of lower baseline serological values, (≤ 30 ng/mL) individually higher values were administered.
- Daily intake of antioxidants (such as a handful of berries, a cup of green tea, coffee without milk, one pinch of Curcuma and/or ginger).
- Daily intake of fibre (vegetables, fruits, legumes, and whole grains).
- Daily intake of nitrate-containing plants (such as a portion of spinach, beetroot or rocket).

Sample collection

Samples were collected at baseline, after 1 month, and after 2 months.

Immunological assessment

Four mL of unstimulated whole saliva (UWS) was collected before mouthwash samples and periodontal examination by using the resting drooling method. Participants were instructed to sit comfortably in an upright position, with their heads slightly down to pool saliva in their mouths, and to allow saliva to fall into 5.0 mL sterile cryotubes.²⁹

The investigation of inflammatory cytokines IL-6, IL-10, and TNF- α was done using commercially available Enzyme-linked Immunosorbent Assay (ELISA) kits.

Microbiological assessment

Mouthwash samples were collected after plaque index recording and before the assessment of probing depth and bleeding on probing. The samples were collected by rinsing with 10 mL sterile saline (0.9% w/v sodium chloride, KSA) for 30 seconds and spitting the same amount in 15 mL sterilized conical tubes (VACUETTE, Austria).³⁰ Real-time PCR was used to quantify the following bacteria (*S. mutans*, *A. actinomycetemcomitans*, *T. forsythia*, *P. gingivalis*, and *F. nucleatum*). After collection, both samples were immediately kept in an icebox and then stored in a -80°C (Electrolux ML 402, Sweden) freezer until analysis.

Follow-up visits

After the baseline clinical examination, sample collection, and anthropometric assessment (T_0), patients underwent the same procedures at the 1-month (T_1) and 2-month (T_2) follow-up visits, as shown in Figure 1.

Methods of assessment

Anthropometric assessment

Participants underwent anthropometric analysis using the InBody-770 scale (InBody Co., Ltd., South Korea). Body composition analysis determined weight, BMI, skeletal muscle mass (SMM), and percent body fat (PBF).

Clinical examination

The first clinical periodontal parameter recorded was the plaque index (PI), according to Silness and L e.³¹ Next, the gingival index (GI) was measured following the method of L e and Silness.³² Bleeding on probing (BOP) was then assessed according to Ainamo and Bay,³³ with all 6 surfaces of each tooth evaluated to determine whether probing elicited bleeding (+) or not (–) within 10 seconds. The severity of BOP was expressed as a percentage. Finally, probing pocket depth (PPD) was measured using standardized periodontal probes (UNC 15, MEDENSY, Italy) at 6 sites per tooth across the entire dentition (mesiobuccal,

midbuccal, distobuccal, mesiolingual, midlingual, and distolingual). One dentist who was calibrated by an expert periodontist conducted the clinical examination for all study participants.

Microbiological assessment


Once the samples were received at the JUST Research Labs, they were transferred and stored in a deep freezer at -70°C before they were analyzed using a real-time polymerase chain reaction (RT-PCR) machine. Immediately before analysis, samples were removed from the deep freezer and DNA in each mouthwash sample was extracted using the DNeasy[®] PowerSoil[®] Pro Kit (Qiagen, Hilden, Germany). The concentration and purity of DNA were evaluated using a Biotek PowerWave XS2 Spectrophotometer (BioTek Instruments, Inc., Winooski, VT, USA).

DNA extraction: The DNeasy kit was used according to the manufacturer's instructions (DNeasy Kit Handbook), allowing rapid and efficient purification of total cellular DNA.

DNA quantification using RT-PCR: For positive control, the plasmid cloning vector (pUC 19) of known DNA concentration was used and the PCR target sequence was inserted into it. The plasmid was diluted in a 7-time serial dilution. Each DNA sample was tested in triplicate; 1 negative control and 2 positive controls of bacterial dilution 10^{-5} ng/ μL and 10^{-6} ng/ μL were used for the RT-PCR reaction.³⁴ The copy number of a plasmid DNA template was determined by using the equation at the website cels.uri.edu/gsc/cndna.html for DNA of *Streptococcus mutans*, *Aggregatibacter actinomycetemcomitans*, *Tannerella forsythia*, *Porphyromonas gingivalis*, and *Fusobacterium nucleatum*.

The machine used for RT-PCR was the Rotor-Gene Q MDx 5 plex system (Qiagen, Hilden, Germany), and the BlasTaq[™] 2X qPCR MasterMix Kit (Applied Biological Materials, Inc., Canada, Cat. No. G891) was used to detect amplified DNA according to manufacturer instructions.

Figure 1. Initial assessment and follow-up visits



For the RT-PCR reaction, a mixture of the following components was prepared: 10 µL of BlasTaq™ 2X qPCR MasterMix, 0.5 µL of Forward Primer (10 µM), 0.5 µL of Reverse Primer (10 µM), and 7 µL of Nuclease-free H₂O. An 18 µL of this mixture was placed in a 1.5 mL strip tube and 2 µL of the extracted DNA was added. Finally 20µL of the reaction mix was obtained. The SYBR Green dye source wavelength was 470 nm and the detection wavelength was 510 nm. Each run in the RT-PCR machine took approximately 100 minutes and was composed of 1 cycle for enzyme activation under 95 °C for 3 min, 40 cycles for denaturation, and annealing/extension under 95 °C, 60 °C for 15 sec, 1 min, respectively. Melting curve obtained under 55 °C to 59 °C to confirm the specificity of primers to the amplified target DNA sequence.

The forward and reverse primers were used for all bacterial species³⁵⁻³⁹ as shown in Table 1.

Immunological assessment

Unstimulated whole saliva (UWS) collected at baseline, after 1 month, and at the end of the study after 2 months was examined using Enzyme-linked Immunosorbent Assay (ELISA) to detect the levels of inflammatory cytokines (IL-6, TNF-α, and IL-10). The kits used by ELK Biotechnology CO., Ltd, China, were Human IL-6 ELISA Kit Cat: ELK1156, Human TNF-α ELISA Kit Cat: ELK1190, and Human IL-10 ELISA Kit Cat: ELK1142. ELISA analysis was performed according to manufacturer instructions for each inflammatory cytokine. The wavelength for ELISA was 450 nm. Samples were analyzed on the Biotek PowerWave XS2 Spectrophotometer (BioTek Instruments, Inc., Winooski, VT, USA).

Statistical analysis

Data were described using means and standard deviations for normally distributed variables. Changes in parameters over time were compared across the 3 timelines using repeated measures ANOVA. Data analysis was performed using SPSS version 25 (IBM Corporation, USA), and a *p* value ≤0.05 was considered statistically significant.

RESULTS

Seventy potential participants were assessed for eligibility. Thirteen refused to participate, and twenty-three did not meet the inclusion criteria. Thirty-four participants were initially included in the study. However, 11 were excluded: 2 took antibiotics, 7 did not comply with sample collection, and 2 missed follow-up visits with the dietitian (Figure 2). Twenty-three participants completed the study and were included in the analysis. The mean age of the participants was 25.7 ± 6.8 years, with 82.6% holding a bachelor's degree and 17.4% a master's degree. Demographic data are shown in Table 2.

Anthropometric outcomes

The anthropometric parameters results were available for 23 patients at 3 timelines. After 2 months, significant reductions were observed in the mean differences of all anthropometric parameters, including weight, BMI, SMM, and PBF. The reductions were -3.8 kg, -1.3 kg/m², -0.5 kg, and -1.6 %, respectively, as shown in Table 3.

Figure 2. Flowchart of participants




Table 1. Forward and reverse RT-PCR primer sequences for the examined bacterial species

Bacterium	Forward sequence	Reverse sequence
<i>S. mutans</i> ²⁷	GCCTACAGCTCAGAGATGCTATTCT	GCCATACACCACTCATGAATTGA
<i>A. actinomycetemcomitans</i> ²⁸	CTTACTACTCTTGACATCCGAA	ATGCAGCACCTGTCTCAAAGC
<i>T. forsythia</i> ²⁹	GGGTGAGTAACGCGTATGTAACCT	ACCCATCCGCAACCAATAAA
<i>P. gingivalis</i> ³⁰	TGGTTTCATGCAGCTTCTTT	TCGGCACCTTCGTAATTCTT
<i>F. nucleatum</i> ³¹	AAGCGCGTCTAGGTGGTTATGT	TGTAGTCCGCTACCTCTCCAG

Table 2. Demographic data

Total number of participants	Age (years)		Educational level (%)	
	Mean \pm SD	Range	Bachelor's degree	Master's degree
23	25.7 \pm 6.8	18 to 45	82.6	17.4

Values are expressed as means and standard deviations

Table 3. Anthropometric parameters at different timelines

Variables	T ₀ (N = 23)		T ₁ (N = 23)		T ₂ (N = 23)		Net difference	P value
	Mean	SD	Mean	SD	Mean	SD		
Weight (kg)	84.6	19.4	81.8	17.7	80.8	17.3	-3.8	0.01
BMI (kg/m ²)	31.9	6.8	30.9	6.3	30.6	6.1	-1.3	0.01
SMM (kg)	25.6	5.5	25.4	5.2	25.1	5.1	-0.5	0.02
PBF (%)	43.9	7.1	42.6	7.3	42.3	7.4	-1.60	0.01

Values are expressed as means and standard deviations. A *p* value \leq 0.05 is considered statistically significant. BMI = body mass index; SMM = skeletal muscle mass; PBF = percent body fat.

Table 4. Clinical parameters at different timelines

Clinical parameters	T ₀ (N = 23)		T ₁ (N = 23)		T ₂ (N = 23)		P value
	Mean	SD	Mean	SD	Mean	SD	
BOP%	37	0.16	24	0.12	19	0.08	0.01
GI	0.77	0.32	0.47	0.24	0.38	0.17	0.01
PI	1.14	0.48	0.69	0.57	0.54	0.38	0.01
PPD	1.83	0.22	1.53	0.29	1.35	0.3	0.01

Values are expressed as means and standard deviations. A *p* value \leq 0.05 is considered statistically significant. BOP = bleeding on probing; GI = gingival index; PI = plaque index; PPD = periodontal probing depth.

Table 5. Average copy number of bacteria (μL) at different timelines

Bacterium	T ₀ (N = 23)		T ₁ (N = 23)		T ₂ (N = 23)		p value
	Mean	SD	Mean	SD	Mean	SD	
<i>S. mutans</i>	10.526	10.292	10.449	10.249	10.484	10.415	0.42
<i>A. actinomycetemcomitans</i>	12.747	12.273	12.719	12.181	12.744	12.208	0.36
<i>F. nucleatum</i>	12.456	11.811	12.479	11.774	12.479	11.786	0.34
<i>T. forsythia</i>	9.566	9.147	9.717	9.456	9.622	9.246	0.01
<i>P. gingivalis</i>	9.274	9.235	9.279	9.184	9.365	9.358	0.42

Values are expressed as means and standard deviations. A p value ≤ 0.05 is considered statistically significant.

Clinical outcomes

After 2 months, significant reductions were observed in the mean differences of all clinical periodontal parameters, including BOP, GI, PI, and PPD, with a p value of 0.01, as shown in Table 4.

Microbiological outcomes

The microbiological results were available for all samples at 3 timelines for the 5 bacterial species. According to the mean bacterial counts at the 3 timelines, the highest count was for *A. actinomycetemcomitans*, while the lowest was for *P. gingivalis*. No significant mean differences for the bacteria were observed after 2 months except for *T. forsythia*, which showed a significant increase in the mean differences with a p value of 0.01, as shown in Table 5.

Immunological outcomes

The results of the ELISA analysis were available for 23 patients at 3 timelines. No significant differences were detected in the mean values of all salivary inflammatory cytokines IL-6, IL-10, and TNF- α after 2 months, as shown in Table 6.

DISCUSSION

The influence of diet on the oral microbial ecosystem and its correlation with oral diseases, such as caries and periodontal disease, has been under focus in the last 2 decades. Although a few studies have investigated the influence of AID on gingivitis in individuals with normal weight,^{12,19–22} there is no known published research on the association between AILCD and gingivitis in women with obesity. The key findings of this study were the significant reductions in clinical and anthropometric parameters. However, there was a significant increase in *T. forsythia* counts after 2 months.

In terms of demographics, this study included 23 female participants, as an association between obesity and periodontal disease has been identified in females, with the link becoming more pronounced as obesity levels increase.^{25,40} Interestingly, this pattern does not appear in males with higher BMI. This discrepancy may be because, in men, a higher BMI often reflects increased muscle mass or bone density rather than excess body fat. Additionally, fat distribution differs between genders, so a higher BMI in

Table 6. Inflammatory cytokines at different timelines

Cytokines	T ₀ (N = 23)		T ₁ (N = 23)		T ₂ (N = 23)		p value
	Mean pg/mL	SD	Mean pg/mL	SD	Mean pg/mL	SD	
IL-6	110.6	36.2	101	34.4	102.5	18.8	0.54
IL-10	63.7	15	78.9	38	62.5	22.8	0.13
TNF- α	125.1	27.5	136.7	32.8	125.7	29	0.33

Values are expressed as means and standard deviations. IL-6 = Interleukin-6; IL-10 = Interleukin-10; TNF- α = Tumour necrosis factor α .

men does not necessarily indicate obesity.²³

The consistent reduction in anthropometric parameters in the present study emphasized the beneficial effect of AILCD in weight loss and obesity management. In agreement with the current study's findings, Woelber et al.²² observed significant weight loss in the test group, attributed to a lower total energy intake from the AID. A recent study showed that a low-carbohydrate diet affects body composition by reducing weight and body fat, and when coupled with physical activity, it helps preserve muscle mass.⁴¹ The significant loss of SMM in this study can be attributed to the 1500-calorie diet with 20% carbohydrates (75 g/day) and 25% protein (94 g/day), which induces caloric restriction. In this case, the body breaks down muscle for energy, leading to further muscle protein breakdown. However, introducing resistance exercises along with the diet helps preserve muscle mass while promoting fat loss.

Regarding the clinical findings of the current study, the participants experienced a reduction in the severity of gingival inflammatory parameters likely because of the restricted sugar intake and the adoption of an anti-inflammatory dietary regimen. The baseline findings support the hypothesis that a modern Western diet, high in refined carbohydrates and with an elevated omega-6 to omega-3 fatty acid ratio, contributes to increased inflammatory processes.⁴²

The reduction in PI can also be attributed to the Hawthorne effect because all participants were aware that they would be examined after brushing their teeth; they may have brushed their teeth more meticulously than usual.⁴³ There is also a hypothesis that a reduction in inflammation leads to a decrease in plaque formation.³ The findings of the current study are in agreement with clinical trial findings where there was a significant reduction in gingival inflammation parameters (PI, GI, BOP, and PPD) even in the absence of interdental hygiene.²² In contrast, Baumgartner et al.¹¹ found that the association of plaque with gingival inflammation might only be valid under Western diet conditions. As PI increased, BOP and PPD decreased significantly in this study. Similar results were observed by El Makaky et al.²¹ in which gingival inflammation and BOP were significantly reduced, but there was an increase in PI in the test and control groups. In a controlled clinical trial, Woelber et al.¹⁹ observed that a diet low in carbohydrates, high in omega-3 fatty acids, high in vitamins C and D, and high in fibre could significantly reduce gingival and periodontal inflammation. Despite constant plaque values in both groups, GI, BOP, and PPD were significantly reduced to roughly half of the baseline values in the experimental group. In a study by Rajaram et al.,¹² significant reductions were observed in the test group compared with the control group in GI and BOP. However, no significant differences were observed in the PI in the test group compared with the control group.

The findings of the previously mentioned studies align with the current study on clinical periodontal parameters (GI, BOP, and PPD). However, there were differences in the PI findings, which supports the crucial debate about the role of plaque as an etiological factor for gingivitis under various dietary conditions, emphasizes its contribution to gingival inflammation in Western diet conditions, and highlights the importance of host modulation.^{7,11,43,44}

Regarding the microbial counts of *S. mutans* and *A. actinomycetemcomitans*, the current study's findings contrast with those of Baumgartner et al.¹¹ in which *Streptococcus* species and *A. actinomycetemcomitans* counts in tongue samples were significantly reduced after 4 weeks of the Stone Age diet compared to baseline. Regarding *F. nucleatum*, the findings of the current study align with those of Woelber et al.²² where there were no significant changes in the counts of *F. nucleatum* after 4 weeks of AID. In contrast, Tennert et al.²⁰ observed a significant reduction in the total count of *Fusobacterium* spp. in plaque samples from a healthy diet group. Conversely, Baumgartner et al.¹¹ found a significant increase in the total count of *F. nucleatum* from subgingival samples after 4 weeks of the Stone Age diet.

The current study revealed a significant increase in *T. forsythia* counts after 2 months. This finding may be related to the BMI (31.9 kg/m² at baseline) of the subjects; these women with obesity may have experienced an overgrowth of *T. forsythia* due to obesity-related dysbiosis in periodontal pathogens. Therefore, preventive measures of longer dietary interventions, physical activity, and periodontal care are recommended to alleviate the complications of obesity-related periodontal dysbiosis. Woelber et al.²² had contradictory results, finding no significant differences in the total counts of subgingival *T. forsythia* after 4 weeks of AID. Yet, Baumgartner et al.¹¹ observed a significant reduction in the total counts of *T. forsythia* from tongue samples in all subjects after 4 weeks of the Stone Age diet.

In the current study, there were no significant differences in the *P. gingivalis* counts after 2 months. Likewise, Woelber et al.²² found no significant differences in the total count of *P. gingivalis* in subjects with gingivitis after 4 weeks of AID.

The difference in periodontal pathogen counts may be due to heterogeneity between studies in terms of the subjects' BMI, adherence to the diet, oral hygiene measures, gingival immunological reactions, and physical activity.

Despite the clinical reduction in the periodontal parameters, no significant changes in the periodontal pathogens were observed, except for *T. forsythia*. These findings require further discussion as they relate to the etiological significance of the dental biofilm. First, it is possible that the clinical outcomes were generated by an altered immunological reaction in the periodontal tissues rather than an altered microbial composition of the dental

biofilm. Second, there could have been direct effects on the periodontal microbiota that DNA sequencing did not detect. The intervention may not have affected the counts of bacteria but rather their metabolic activity.⁴⁵

Regarding the immunological findings of the salivary cytokines IL-6, IL-10, and TNF- α , there were no significant differences in all cytokines after 2 months. A possible explanation for this is that obese individuals have elevated levels of salivary IL-6 and TNF- α , which are used as an alternative method to serum TNF- α in diagnosing obesity. Additionally, many of the participants were young adults (25.7 ± 6.8 years) and the salivary cytokine concentration declines with age. This correlation may be due to different age-related changes, such as a decrease in total salivary protein concentration and structural and histological variations in salivary glands, such as acinar cell atrophy and replacement of normal parenchyma with fibrosis. Third, the sampling technique by collecting unstimulated whole saliva (UWS) may affect the concentration of salivary cytokines. UWS has a lower salivary flow rate and a higher possibility of sample blood contamination compared with stimulated whole saliva. These factors could alter the salivary cytokine concentration in UWS samples.

The current study's findings align with those of Woelber et al.²² who showed no significant changes in serum IL-6 and TNF- α after 4 weeks of AID. Moreover, Ruth et al.⁴⁵ observed that serum TNF- α level did not differ after 12 weeks of a high-fat, low-carbohydrate diet. In contrast, El Makaky et al.²¹ found a significant reduction in salivary TNF- α and IL-6 levels within a test group after 4 weeks of AID in children with gingivitis. These contradictory results may be due to differences in the sampling techniques and the kits used for the assessments in the studies.

Although clinical periodontal parameters were reduced during the study, periodontal pathogens and inflammatory cytokines did not change significantly in these women with obesity. More time may be required to achieve greater weight loss and, subsequently, a reduction in periodontal pathogens.

Study limitations

The main limitation of the current study is the absence of a control group, a relatively small sample size, and a short follow-up period. Future studies are recommended to conduct long-term follow-up RCTs with larger sample sizes, non-obese periodontitis patients, and a broader range of bacterial species. Furthermore, developing a protocol to ensure participant adherence to the diet, such as a food delivery service for diet standardization, is advisable.

CONCLUSION

The AILCD used in this study was associated with significant improvements in clinical periodontal and anthropometric parameters. However, *Tannerella forsythia* showed a significant increase in women with obesity.

ACKNOWLEDGEMENTS

The authors thank Jordan University of Science and Technology, Irbid-Jordan, for funding this research (Grant number: 20220426).

CONFLICTS OF INTEREST

The authors declare that they have no conflicts of interest.

REFERENCES

1. Trombelli L, Farina R, Silva CO, Tatakis DN. Plaque-induced gingivitis: case definition and diagnostic considerations. *J Clin Periodontol.* 2018;45(Suppl 20):S44–S67.
2. Murakami S, Mealey BL, Mariotti A, Chapple ILJ. Dental plaque-induced gingival conditions. *J Clin Periodontol.* 2018;45:S17–S27.
3. Van Der Velden UJ. The significance of supragingival plaque accumulation in periodontal disease. *Int J Dent Hyg.* 2006;4(Suppl 1):11–14.
4. Lang NP, Schätzle MA, Löe HJ. Gingivitis as a risk factor in periodontal disease. *J Clin Periodontol.* 2009;36:3–8.
5. Stöhr J, Barbaresco J, Neuenschwander M, Schlesinger S. Bidirectional association between periodontal disease and diabetes mellitus: a systematic review and meta-analysis of cohort studies. *Sci Rep.* 2021;11(1):13686.
6. Herrera D, Sanz M, Shapira L, Brotons C, Chapple I, Frese T, et al. Association between periodontal diseases and cardiovascular diseases, diabetes and respiratory diseases: Consensus report of the Joint Workshop by the European Federation of Periodontology (EFP) and the European arm of the World Organization of Family Doctors (WONCA Europe). *J Clin Periodontol.* 2023;50(6):819–41.
7. Woelber JP, Bartha V, Baumgartner S, Tennert C, Schlagenhauf U, Ratka-Krüger P, et al. Is diet a determining factor in the induction of gingival inflammation by dental plaque? A secondary analysis of clinical studies. *Nutrients.* 2024;16(7):923.
8. Loesche WJJ. Role of *Streptococcus mutans* in human dental decay. *Microbiol Rev.* 1986;50(4):353–80.
9. Gasmi Benahmed A, Gasmi A, Dadar M, Arshad M, Björklund G. The role of sugar-rich diet and salivary proteins in dental plaque formation and oral health. *J Oral Biosci.* 2021;63(2):134–41.
10. Hajjshengallis G, Chavakis T, Lambris JD. Current understanding of periodontal disease pathogenesis and targets for host-modulation therapy. *Periodontology 2000.* 2020;84(1):14–34.
11. Baumgartner S, Imfeld T, Schicht O, Rath C, Persson RE, Persson GR. The impact of the stone age diet on gingival conditions in the absence of oral hygiene. *J Periodontol.* 2009;80(5):759–68.
12. Rajaram SS, Nisha S, Ali NM, Shashikumar P, Karmakar S, Pandey VJ. Influence of a low-carbohydrate and rich in Omega-3 fatty acids, ascorbic acid, antioxidants, and fiber diet on clinical outcomes in patients with chronic gingivitis: a randomized controlled trial. *J Int Soc Prev Community Dent.* 2021;11(1):58–67.
13. Martinon P, Fraticelli L, Giboreau A, Dussart C, Bourgeois D, Carrouel FJ. Nutrition as a key modifiable factor for periodontitis and main chronic diseases. *J Clin Med.* 2021;10(2):197.

14. Santonocito S, Polizzi A, Palazzo G, Indelicato F, Isola G. Dietary factors affecting the prevalence and impact of periodontal disease. *Clin Cosmet Investig Dent*. 2021;13:283–92.
15. Calder PC. Omega-3 fatty acids and inflammatory processes: from molecules to man. *Biochem Soc Trans*. 2017;45(5):1105–1115.
16. Wasti J, Wasti A, Singh RJ. Efficacy of antioxidants therapy on progression of periodontal disease—a randomized control trial. *Indian J Dent Res*. 2021;32(2):187–91.
17. Fu Z, Gilbert ER, Liu D. Regulation of insulin synthesis and secretion and pancreatic Beta-cell dysfunction in diabetes. *Curr Diabetes Rev*. 2013;9(1):25–53.
18. Atkinson FS, Khan JH, Brand-Miller JC, Eberhard J. The impact of carbohydrate quality on dental plaque pH: Does the glycemic index of starchy foods matter for dental health? *Nutrients*. 2021;13(8):2711.
19. Woelber JP, Bremer K, Vach K, König D, Hellwig E, Ratka-Krüger P, et al. An oral health optimized diet can reduce gingival and periodontal inflammation in humans—a randomized controlled pilot study. *BMC Oral Health*. 2017;17(1):1–8.
20. Tennert C, Reinmuth AC, Bremer K, Al-Ahmad A, Karygianni L, Hellwig E, et al. An oral health optimized diet reduces the load of potential cariogenic and periodontal bacterial species in the supragingival oral plaque: a randomized controlled pilot study. *Microbiologyopen*. 2020;9(8):e1056.
21. El Makaky YM, Beltagy TM, El Makakey AM. The effects of an anti-inflammatory diet on gingival health in children: randomized controlled trial. *Egyptian Dent J*. 2019;65(3):1995–2002.
22. Woelber JP, Gärtner M, Breuninger L, Anderson A, König D, Hellwig E, et al. The influence of an anti-inflammatory diet on gingivitis. A randomized controlled trial. *J Clin Periodontol*. 2019;46(4):481–90.
23. Dalla Vecchia CF, Susin C, Rösing CK, Oppermann RV, Albandar MJM. Overweight and obesity as risk indicators for periodontitis in adults. *J Periodontol*. 2005;76(10):1721–1728.
24. Haffajee AD, Socransky SS. Relation of body mass index, periodontitis and *Tannerella forsythia*. *J Clin Periodontol*. 2009;36(2):89–99.
25. Suvan JE, Finer N, D'Aiuto FJP. Periodontal complications with obesity. *Periodontology 2000*. 2018;78(1):98–128.
26. Gaio EJ, Haas AN, Rösing CK, Oppermann RV, Albandar JM, Susin C. Effect of obesity on periodontal attachment loss progression: a 5-year population-based prospective study. *J Clin Periodontol*. 2016;43(7):557–65.
27. Chapple IL, Mealey BL, Van Dyke TE, Bartold PM, Dommisch H, Eickholz P, et al. Periodontal health and gingival diseases and conditions on an intact and a reduced periodontium: Consensus report of workgroup 1 of the 2017 World Workshop on the Classification of Periodontal and Peri-Implant Diseases and Conditions. *J Periodontol*. 2018;89:S74–S84.
28. Feinman RD, Pogozelski WK, Astrup A, Bernstein RK, Fine EJ, Westman EC, et al. Dietary carbohydrate restriction as the first approach in diabetes management: critical review and evidence base. *Nutrition*. 2015;31(1):1–13.
29. Kong X, Ferracane R, De Luca L, Vitaglione P. Salivary concentration of N-acylethanolamines upon food mastication and after meal consumption: Influence of food dietary fiber. *Food Res Int*. 2016;89(Pt 1):186–93.
30. Boutaga K, Savelkoul PH, Winkel EG, van Winkelhoff AJ. Comparison of subgingival bacterial sampling with oral lavage for detection and quantification of periodontal pathogens by real-time polymerase chain reaction. *J Periodontol*. 2007;78(1):79–86.
31. Silness J, Loe H. Periodontal disease in pregnancy II. Correlation between oral hygiene and periodontal condition. *Acta Odontol Scand*. 1964;22(1):121–35.
32. Loe H, Silness J. Periodontal disease in pregnancy I. Prevalence and severity. *Acta Odontol Scand*. 1963;21(6):533–51.
33. Ainamo J, Bay I. Problems and proposals for recording gingivitis and plaque. *Int Dent J*. 1975;25(4):229–35.
34. Mikić A, Alomari A, Gowers DM. Classical recombinant DNA cloning. *Methods Mol Biol*. 2023;2633:1–24.
35. Buonavoglia A, Trotta A, Camero M, Cordisco M, Dimuccio MM, Corrente M. Streptococcus mutans associated with endo-periodontal lesions in intact teeth. *Appl Sci*. 2022;12(22):11837.
36. Kure K, Sato H, Aoyama N, Izumi Y. Accelerated inflammation in peripheral artery disease patients with periodontitis. *J Periodontal Implant Sci*. 2018;48(6):337–46.
37. Xu Y, Selerio-Poely T, Ye X. Clinical and microbiological effects of egg yolk antibody against Porphyromonas gingivalis as an adjunct in the treatment of moderate to severe chronic periodontitis: a randomized placebo-controlled clinical trial. *J Periodontal Implant Sci*. 2018;48(1):47–59.
38. Barbosa GM, Colombo AV, Rodrigues PH, Simionato MRL. Correction: Intraspecies variability affects heterotypic biofilms of Porphyromonas gingivalis and Prevotella intermedia: Evidences of strain-dependence biofilm modulation by physical contact and by released soluble factors. *PLoS One*. 2015;10(11):e0143903.
39. Tadokoro K, Yamaguchi T, Kawamura K, Shimizu H, Egashira T, Minabe M, et al. Rapid quantification of periodontitis-related bacteria using a novel modification of Invader PLUS technologies. *Microbiol Res*. 2010;165(1):43–49.
40. de Moraes GBD, de Souza VSM, da Silva Mazzeti CM, Macedo MLR, Rafacho BPM. The effects of the low-carb diet in the body composition and weight loss. *Nutrire*. 2024;49(1):11.
41. Bosma-den Boer MM, van Wetten ML, Pruimboom L. Chronic inflammatory diseases are stimulated by current lifestyle: How diet, stress levels and medication prevent our body from recovering. *Nutr Metab (Lond)*. 2012;9(1):32.
42. McCarney R, Warner J, Iliffe S, Van Haselen R, Griffin M, Fisher P. The Hawthorne Effect: a randomised, controlled trial. *BMC Med Res Methodol*. 2007;7(1):1–8.
43. Bartha V, Exner L, Schweikert D, Woelber JP, Vach K, Meyer AL, et al. Effect of the Mediterranean diet on gingivitis: a randomized controlled trial. *J Clin Periodontol*. 2022;49(2):111–22.
44. Albenberg LG, Wu GD. Diet and the intestinal microbiome: Associations, functions, and implications for health and disease. *Gastroenterology*. 2014;146(6):1564–1572.
45. Ruth MR, Port AM, Shah M, Bourland AC, Istfan NW, Nelson KP, et al. Consuming a hypocaloric high fat low carbohydrate diet for 12 weeks lowers C-reactive protein, and raises serum adiponectin and high density lipoprotein-cholesterol in obese subjects. *Metabolism*. 2013;62(12):1779–1787.